

Theory of internal transitions of charged excitons in quantum wells in magnetic fields

A. B. Dzyubenko

*Institut für Theoretische Physik, J.W. Goethe-Universität, 60054 Frankfurt, Germany
General Physics Institute, RAS, Moscow 117942, Russia*

A. Yu. Sivachenko

*The Weizmann Institute of Science, Rehovot 76100, Israel
(February 1, 2008)*

For charged semiconductor complexes in magnetic fields B , we discuss an exact classification of states, which is based on magnetic translations. In this scheme, in addition to the total orbital angular momentum projection M_z and electron and hole spins S_e , S_h , a new exact quantum number appears. This oscillator quantum number, k , is related physically to the center of the cyclotron motion of the complex as a whole. In the dipole approximation k is strictly conserved in magneto-optical transitions. We discuss implications of this new exact selection rule for internal intraband magneto-optical transitions of charged excitons X^- in quantum wells in B .

73.20.Dx, 71.70.Di, 76.40.+b

Recently, there has been considerable experimental and theoretical interest in charged excitons X^- and X^+ in magnetic fields B in 2D systems. Experimentally, magneto-optical interband [1] transitions of charged excitons have been studied extensively. Theoretically, the binding of charged excitons X^- has been considered in quantum dots [2], in a strictly-2D system in the high-magnetic field limit [3], and in realistic quantum wells at finite B [4]. In all these theoretical works on charged excitons, the existing exact symmetry — magnetic translations — has not been identified. The aim of the present theoretical work is to describe in some detail this symmetry and its manifestations in intraband magneto-optical transitions (see also [5]). Experimental evidences for internal X^- singlet and triplet transitions have been reported very recently in [6], where also comparison with quantitative calculations is presented.

We consider a system of interacting particles of charges e_j in a magnetic field $\mathbf{B} = (0, 0, B)$ described by the Hamiltonian

$$H = \sum_j \frac{\hat{\pi}_j^2}{2m_j} + \frac{1}{2} \sum_{i \neq j} U_{ij}(\mathbf{r}_i - \mathbf{r}_j), \quad (1)$$

here $\hat{\pi}_j = -i\hbar \nabla_j - \frac{e_j}{c} \mathbf{A}(\mathbf{r}_j)$ is the kinematic momentum operator of the j -th particle in $\mathbf{B}$ and U_{ij} are the potentials of interactions that can be rather arbitrary. Dynamical symmetries of (1) are the following. In the symmetric gauge $\mathbf{A} = \frac{1}{2} \mathbf{B} \times \mathbf{r}$, there is the axial symmetry about the z -axis $[H, \hat{L}_z] = 0$, where $\hat{L}_z = \sum_j (\mathbf{r}_j \times -i\hbar \nabla_j)_z$.

Therefore, the total angular momentum projection M_z , an eigenvalue of $\hat{L}_z$, is a good quantum number. In a uniform $\mathbf{B}$, the Hamiltonian (1) is also invariant under a group of magnetic translations whose generators are the components of the operator $\hat{\mathbf{K}} = \sum_j \hat{\mathbf{K}}_j$, where $\hat{\mathbf{K}}_j = \hat{\pi}_j - \frac{e_j}{c} \mathbf{r}_j \times \mathbf{B}$ (see, e.g., [7]). $\hat{\mathbf{K}}$ is the exact integral of the motion: $[H, \hat{\mathbf{K}}] = 0$. The components of $\hat{\mathbf{K}}$ and $\hat{\pi} = \sum_j \hat{\pi}_j$ commute in $\mathbf{B}$ as

$$[\hat{K}_x, \hat{K}_y] = -[\hat{\pi}_x, \hat{\pi}_y] = -i \frac{\hbar B}{c} Q, \quad Q \equiv \sum_j e_j, \quad (2)$$

while $[\hat{K}_p, \hat{\pi}_q] = 0$, $p, q = x, y$. For neutral complexes (atoms, excitons, biexcitons) $Q = 0$, and classification of states in B are due to the continuous two-component vector — the 2D magnetic momentum $\mathbf{K} = (K_x, K_y)$. For charged systems the components of $\hat{\mathbf{K}}$ cannot be observed simultaneously. This determines the macroscopic Landau degeneracy of exact eigenstates of (1). For a dimensionless operator $\hat{\mathbf{k}} = \sqrt{c/2\hbar B|Q|} \hat{\mathbf{K}}$ we have $[\hat{k}_x, \hat{k}_y] = -iQ/|Q|$. Therefore, $\hat{k}_{\pm} = (\hat{k}_x \pm i\hat{k}_y)/\sqrt{2}$ are Bose raising and lowering ladder operators: $[\hat{k}_+, \hat{k}_-] = -Q/|Q|$. It follows then that $\hat{\mathbf{k}}^2 = \hat{k}_+ \hat{k}_- + \hat{k}_- \hat{k}_+$ has discrete oscillator eigenvalues $2k+1$, $k = 0, 1, \dots$. Since $[\hat{\mathbf{k}}^2, H] = 0$ and $[\hat{\mathbf{k}}^2, \hat{L}_z] = 0$, the exact charged eigenstates of (1), in addition to the electron S_e and hole S_h spin quantum numbers, can be simultaneously labeled by the discrete quantum numbers k and M_z . The labelling therefore is $|k M_z S_e S_h \nu\rangle$. Here ν is the “principal” quantum number, which can be discrete (bound states) or continuous (unbound states forming a continuum) [5]. The $k = 0$ states are *Parent States* (PS’s) within a degenerate manifold. All other *Daughter* states in each ν -th family are generated out of the PS iteratively: for $Q < 0$ $|k, M_z - k, S_e S_h \nu\rangle = (\hat{k}_-)^k |0, M_z, S_e S_h \nu\rangle / \sqrt{k!}$.

Let us discuss now magneto-optical transitions of charged complexes. In the dipole approximation the photon momentum is negligibly small. Therefore, the quantum number k should be conserved in intra- and interband magneto-optical transitions. For interband transitions in a translationally-invariant system with a simple valence band this leads to a striking result [5] that the ground triplet X_t^- ($S_e = 1$) state is dark in photolumines-




FIG. 1. Schematic drawing of bound and scattering electron triplet $2e-h$ states in the lowest LL's $(N_e N_h) = (00), (10)$. The states are labeled by the total angular momentum projection M_z and oscillator quantum number k . Separation between the 2D e^- and h^- layers $d = 0$ (see inset to Fig. 2). The energy $E_0 = \sqrt{\pi/2} e^2 / \epsilon l_B$ parametrizes the 2D system in the limit of high B . Large (small) dots correspond to the bound parent $k = 0$ (daughter $k = 1, 2, \dots$) X^- states; see text for further explanations.

cence at *finite* magnetic fields B and in a quasi-2D system with a possible $e-h$ asymmetry (cf. [3]). Also, conservation of k prohibits shake-up processes of an isolated singlet X_s^- ($S_e = 0$) state in B . In this work we concentrate on internal intraband transitions of charged complexes in a magnetic field. In these transitions conservation of k follows from the commutativity $[\hat{V}^\pm, \hat{\mathbf{K}}] = 0$, where $\hat{V}^\pm \sim \sum_j e_j \pi_j^\pm / m_j$, is the Hamiltonian of the interaction with the radiation of polarization $\sigma^\pm$ and $\pi_j^\pm = \pi_{jx} \pm i\pi_{jy}$. Other selection rules in this case are conservation of spins S_e, S_h and $\Delta M_z = \pm 1$ in the $\sigma^\pm$ polarization for the envelope function. Conservation of k constitutes a new exact selection rule. It leads, in particular, to some striking consequences for bound-to-bound transitions from the triplet X_t^- ground state.

To show this we shall consider the system of strictly-2D e^- and h^- layers separated by a distance d (see inset to Fig. 2) in the limit of high magnetic fields. To understand internal X^- transitions, it is necessary to consider the eigenstates associated with higher Landau levels (LL's). To this end we use expansion in free LL's, which has been described in some detail elsewhere [8,5]. The calculated three-particle $2e-h$ eigenspectra (electrons in the triplet




FIG. 2. Energies and dipole matrix elements of the inter-LL transitions from the ground X_{t00}^- state in the high-field limit. Shown are the cases of three different separations $d = 0, 0.3l_B$, and $0.7l_B$ between the strictly-2D e^- and h^- layers (see inset). A filled dot shows the position of the forbidden bound-to-bound $X_{t00}^- \rightarrow X_{t10}^-$ transition for $d = 0$.

state) in the two lowest LL's are shown for $d = 0$ in Fig. 1. Generally, the eigenspectra associated with each LL consist of bands of *finite* width $\sim E_0 = \sqrt{\pi/2} e^2 / \epsilon l_B$, where $l_B = (\hbar c / eB)^{1/2}$. The states within each such band form a *continuum* corresponding to the extended motion of a neutral magnetoexciton (MX) as a whole with the second electron in a scattering state. As an example, the continuum in the lowest $(N_e N_h) = (00)$ LL consists of the MX band of width E_0 extending down in energy from the free (00) LL. This corresponds to the $1s$ MX ($N_e = N_h = 0$) [9] plus a scattered electron in the zero LL, labeled $X_{00} + e_0$. The structure of the continuum in the $(N_e N_h) = (10)$ LL is more complicated: in addition to the $X_{00} + e_1$ band of the width E_0 , there is another MX band of width $0.574E_0$ also extending down in energy from the free $(N_e N_h) = (10)$ LL. This corresponds to the $2p^+$ exciton ($N_e = 1, N_h = 0$) [9] plus a scattered electron in the $N_e = 0$ LL, labeled $X_{10} + e_0$. There are also bands (not shown in Fig. 1) above each free LL originating from the bound internal motion of two electrons in the absence of a hole [5]. Internal transitions to such bands have extremely small oscillator strengths and not discussed here. Bound X^- states (finite internal motions of all three particles) lie *outside* the continua (Fig. 1). In the limit of high B the only bound X^- state in the zeroth LL $(N_e N_h) = (00)$ is the X^- -triplet. There are no bound X^- -singlet states [3,4] in contrast to the $B = 0$ case. The X^- -triplet binding energy in zero LL's $(N_e N_h) = (00)$ is $0.043E_0$ [3,4]. In the next electron LL $(N_e N_h) = (10)$ there are no bound X^- -singlets, and only one bound triplet state X_{t10}^- , lying below the lower edge

of the MX band [5]. The X_{t10}^- binding energy is $0.086E_0$, twice that of the X_{t00}^- , and similar to the stronger binding of the D^- -triplet in the $N_e = 1$ LL [8].

We focus here on internal transitions in the σ^+ polarization governed by the usual selection rules: spin conserved, $\Delta M_z = 1$. In this case the e -CR-like inter-LL ($\Delta N_e = 1$) transitions are strong and gain strength with B . Both bound-to-bound $X_{t00}^- \rightarrow X_{t10}^-$ and photoionizing X^- transitions are possible. For the latter the final three-particle states in the (10) LL belong to the continuum (Fig. 1), and calculations show that the FIR absorption spectra reflect its rich structure [5]. Transitions to the $X_{00} + e_1$ continuum are dominated by a sharp onset at the edge (transition 1) at an energy $\hbar\omega_{ce}$ plus the X_{t00}^- binding energy. In addition, there is a broader and weaker peak corresponding to the transition to the $X_{01} + e_0$ MX band, transition 2. The latter may be thought of as the $1s \rightarrow 2p^+$ internal transition of the MX [10], which is shifted and broadened by the presence of the second electron. In accordance with this picture, it is visible from Fig. 2 that with increasing separation d between the e - and h -layers (when the exciton binding and, thus, transition energies are reduced and the X - e interaction is effectively diminished), the second peak is redshifted and sharpened. Thus the X^- -triplet behaves physically in the photoionizing bound-to-continuum transitions as an exciton that very loosely binds an electron, and the two “parts” of the complex can absorb the FIR photon, to some extent, independently. The double-peak structure of the bound-to-continuum transitions is a generic feature for transitions from both the singlet and triplet ground X^- states in quasi-2D systems in strong B . Such transitions in translationally invariant systems are discussed theoretically in [6], where also experimental results for bound-to-continuum transitions are reported and comparison between theory and experiment is made.

The inter-LL bound-to-bound transition, $X_{t00}^- \rightarrow X_{t10}^-$, has a very specific spectral position: since the final state is more strongly bound, it lies *below* the e -CR energy $\hbar\omega_{ce} = \hbar eB/m_e c$. However, it has exactly *zero* oscillator strength, a manifestation of the magnetic translational invariance: the two selection rules – conservation of k and $\Delta M_z = 1$ cannot be satisfied simultaneously. Indeed, e.g., the X_{t00} PS (with $k = 0$) has $M_z = -1$, while the X_{t10} PS has $M_z = 1$, so that the usual selection rule $\Delta M_z = 1$ cannot be satisfied. Localization of charged excitons breaks translational invariance and relaxes the k -conservation rule. As a result, the bound-to-bound $X_{t00}^- \rightarrow X_{t10}^-$ transition, which is prohibited in translationally-invariant systems, develops *below* the e -CR [5]. Such a peak is a tell-tale mark of localization of charged triplet excitons.

In conclusion, we have studied the exact symmetry — magnetic translations — for charged excitons in B and established its consequences for intraband magneto-

optical transitions. In particular, we have shown that in translationally invariant quasi-2D system with a simple valence band the bound-to-bound transition from the triplet ground state X_t^- to the next electron Landau level is prohibited. In the presence of translationally-breaking effects (disorder, impurities etc.) the intraband bound-to-bound triplet transition develops below the electron cyclotron resonance. This suggests a method of studying localization of charged excitons.

ABD is grateful to the Alexander von Humboldt Foundation for research support.

-
- [1] See, e.g., K. Kheng, R.T. Cox, Merle Y. d’Aubigné *et al.* Phys. Rev. Lett. **71**, 1752 (1993); G. Finkelstein, H. Shtrikman, and I. Bar-Joseph, Phys. Rev. Lett. **74**, 976 (1995); A.J. Shields, M. Pepper, M.Y. Simmons, and D.A. Ritchie, Phys. Rev. B **52**, 7841 (1995); D. Gekhtman, E. Cohen, Arza Ron, and L.N. Pfeiffer, Phys. Rev. B **54**, 10 320 (1996), D.R. Yakovlev, V.P. Kochereshko, R. Suris *et al.* Phys. Rev. Lett. **79**, 3974 (1997) and references therein.
 - [2] A. Wojs and P. Hawrylak, Phys. Rev. B **51**, 10 880 (1995).
 - [3] J.J. Palacios, D. Yoshioka, and A.H. MacDonald, Phys. Rev. B **54**, R2296 (1996).
 - [4] D.M. Whittaker and A.J. Shields, Phys. Rev. B **56**, 15 185 (1997).
 - [5] A.B. Dzyubenko and A.Yu. Sivachenko, cond-mat/9902086; cond-mat/9908406, to be published in Physica E.
 - [6] A.B. Dzyubenko, A.Yu. Sivachenko, H.A. Nickel, T.M. Yeo, G. Kioussoglou, B.D. McCombe, and A. Petrou, cond-mat/9907124, to be published in Physica E.
 - [7] J.E. Avron, I.W. Herbst, and B. Simon, Annals of Phys. **114**, 431 (1978).
 - [8] A.B. Dzyubenko, Phys. Lett. A **165**, 357 (1992); **173**, 311 (1993).
 - [9] I.V. Lerner and Yu.E. Lozovik, Sov. Phys. JETP **51**, 588 (1980).
 - [10] A.B. Dzyubenko, Pis'ma ZhETF **66**, 588 (1997) [JETP Lett. **66**, 617 (1997)].